

Show *all* of your work and *explain* your answers fully. There is a total of 100 possible points.

As justification for your work, you may use Sage routines designed for matrices, but not routines designed for linear transformations.

1. Consider the following linear transformation from the vector space of polynomials with degree at most 2 to the vector space of 1×2 matrices. (30 points)

$$T: P_2 \rightarrow M_{12}, \quad T(a + bx + cx^2) = [2a - 3b + c \quad a - b + c]$$

- (a) Find the matrix representation relative to the bases $B = \{1 + x, 2x + x^2, -4 + 5x + 4x^2\}$ and $C = \left\{ \begin{bmatrix} 4 & 5 \end{bmatrix}, \begin{bmatrix} 3 & 4 \end{bmatrix} \right\}$.

- (b) Use a version of the Fundamental Theorem of Matrix Representation and your answer to part (a) to compute $T(2 + x + 3x^2)$.

- (c) Find the kernel of T , $\mathcal{K}(T)$, using your answer in part (a).



2. Consider the linear transformation, S . (25 points)

$$S: M_{22} \rightarrow P_3,$$

$$T \left(\begin{bmatrix} a & b \\ c & d \end{bmatrix} \right) = (9a - 2b - 7c - 5d) + (5a - 4b - 9c - 4d)x + (-3a + b + 3c + 2d)x^2 + (-2a + c + d)x^3$$

(a) Build a matrix representation of S and from this determine *as efficiently as possible* that S is invertible.

(b) Find an expression for the outputs of the inverse linear transformation S^{-1} . (In other words, find a “formula” or “rule” for the inverse linear transformation.)

3. Let S_{22} be the vector space of 2×2 symmetric matrices. Provide a basis for S_{22} so that the linear transformation R had a diagonal matrix representation. (20 points)

$$R: S_{22} \rightarrow S_{22}, \quad R \left(\begin{bmatrix} a & b \\ b & c \end{bmatrix} \right) = \begin{bmatrix} 2a - 5b + 5c & 5a - 8b + 5c \\ 5a - 8b + 5c & 5a - 5b + 2c \end{bmatrix}$$



4. Consider the linear transformation from the first problem, T , reprinted below, along with the prior bases $B = \{1 + x, 2x + x^2, -4 + 5x + 4x^2\}$ and $C = \{\begin{bmatrix} 4 & 5 \end{bmatrix}, \begin{bmatrix} 3 & 4 \end{bmatrix}\}$. (25 points)

$$T: P_2 \rightarrow M_{12}, \quad T(a + bx + cx^2) = \begin{bmatrix} 2a - 3b + c & a - b + c \end{bmatrix}$$

- (a) Build a matrix representation of T relative to standard bases (denote these bases as X for the domain and Y for the codomain).

- (b) Build the matrices that relate the representation from part (a) of Problem 1 to the representation in part (a) of this problem, and do the computation that converts one representation into the other.

