Math 290	
Exam 5	
Chapters D, E	

Name:

Show *all* of your work and *explain* your answers fully. There is a total of 100 possible points. For computational problems, place your answer in the provided boxes. Partial credit is proportional to the quality of your explanation. Unless the problem directions specify otherwise, you may use Sage to manipulate matrices and vectors, row-reduce matrices, compute determinants, compute and factor characteristic polynomials, and compute eigenvalues and eigenspaces. No other use of Sage may be used as justification for your answers. When you use Sage be sure to explain your input and show any relevant output (rather than just describing salient features of something I can't see).

1. Compute the determinant of the following matrix by hand. Sage may not be used to justify your answer and you must show enough work to demonstrate that you know how determinants are computed. (10 points)

	$\left\lceil -2 \right\rceil$	-1	3]	
A =	3	1	-2	
	3	1	-1	

nswer:	

2. Suppose that A is an 8×8 matrix with det(A) = 5. Determine, with explanation, det(2A). (15 points)

Answer:		

3. Consider the matrix C. Use Sage only as directed in your justifications of your answers. (40 points)

$$C = \begin{bmatrix} -5 & -2 & -6 & 2\\ 0 & -13 & 16 & 16\\ 4 & 1 & 6 & -1\\ -4 & -13 & 9 & 16 \end{bmatrix}$$

(a) Compute the eigenvalues, and algebraic multiplicities, of C. You may use Sage to find a factored version of the characteristic polynomial.

Answer:			

(b) Compute the eigenspaces, and geometric multiplicities, of C. You may use Sage to row-reduce matrices.

Answer:		

(c) Is C diagonalizable? Why or why not?

4. Are the matrices E and F similar? Why or why not? (15 points)

$$E = \begin{bmatrix} 7 & 3 & 3 \\ -6 & -1 & -6 \\ -2 & -1 & 0 \end{bmatrix} \qquad \qquad F = \begin{bmatrix} 2 & 1 & 1 \\ 3 & 2 & 3 \\ -3 & -1 & -2 \end{bmatrix}$$

5. Construct a 3×3 matrix whose eigenvalues are 1, 2 and 5, and it is not obvious by appearance that these are the eigenvalues of the matrix. Provide enough of a description of your procedure to convince the reader that you have only used the allowed procedures from Sage. (15 points)

6. For a matrix A of size n and a scalar $\alpha \in \mathbf{C}$, determine a formula for det (αA) in terms of α and det(A), and give a proof of your formula. Hint: review your work in the second question. (15 points)